

LA-UR-17-27868

Approved for public release; distribution is unlimited.

Title: Neutron Electric Dipole Moment

Author(s): Bhattacharya, Tanmoy

Intended for: Lattice QCD, 2017-08-28 (Santa Fe, New Mexico, United States)

Issued: 2017-08-31

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Neutron Electric Dipole Moment

Tanmoy Bhattacharya

Los Alamos National Laboratory

August 30, 2017

Introduction

Standard Model CP Violation

Two sources of CP violation in the Standard Model.

- Complex phase in CKM quark mixing matrix.
 - Too small to explain baryon asymmetry
 - Gives a tiny ($\sim 10^{-32}$ e-cm) contribution to nEDM

Dar arXiv:hep-ph/0008248.

- CP-violating mass term and effective $\Theta G\tilde{G}$ interaction related to QCD instantons
 - Effects suppressed at high energies
 - nEDM limits constrain $\Theta \lesssim 10^{-10}$


Crewther *et al.*, *Phys. Lett. B* **88** (1979) 123.

Contributions from beyond the standard model

- Needed to explain baryogenesis
- May have large contribution to EDM

Introduction

Effective Field Theory



Introduction

BSM Operators

Standard model CP violation in the weak sector.

Strong CP violation from dimension 3 and 4 operators anomalously small.

- Dimension 3 and 4:
 - CP violating mass $\bar{\psi}\gamma_5\psi$.
 - Topological charge $G_{\mu\nu}\tilde{G}^{\mu\nu}$.
- Suppressed by v_{EW}/M_{BSM}^2 :
 - Electric Dipole Moment $\bar{\psi}\Sigma_{\mu\nu}\tilde{F}^{\mu\nu}\psi$.
 - Chromo Dipole Moment $\bar{\psi}\Sigma_{\mu\nu}\tilde{G}^{\mu\nu}\psi$.
- Suppressed by $1/M_{BSM}^2$:
 - Weinberg operator (Gluon chromo-electric moment):
 $G_{\mu\nu}G_{\lambda\nu}\tilde{G}_{\mu\lambda}$.
 - Various four-fermi operators.

Introduction

Form Factors

Vector form-factors

Dirac F_1 , Pauli F_2 , **Electric dipole F_3** , and **Anapole F_A**

Sachs electric $G_E \equiv F_1 - (q^2/4M^2)F_2$ and magnetic $G_M \equiv F_1 + F_2$

$$\begin{aligned} \langle N | V_\mu(q) | N \rangle = & \bar{u}_N \left[\gamma_\mu F_1(q^2) + i \frac{[\gamma_\mu, \gamma_\nu]}{2} q_\nu \frac{F_2(q^2)}{2m_N} \right. \\ & + (2i m_N \gamma_5 q_\mu - \gamma_\mu \gamma_5 q^2) \frac{F_A(q^2)}{m_N^2} \\ & \left. + \frac{[\gamma_\mu, \gamma_\nu]}{2} q_\nu \gamma_5 \frac{F_3(q^2)}{2m_N} \right] u_N \end{aligned}$$

- The charge $G_E(0) = F_1(0) = 0$.
- $G_M(0)/2M_N = F_2(0)/2M_N$ is the (anomalous) magnetic dipole moment.
- $F_3(0)/2m_N$ is the electric dipole moment.
- F_A and F_3 violate P; F_3 violates CP.

Dirac Equation

Free neutrons

By Lorentz invariance, an on-shell spin- $\frac{1}{2}$ massive field ψ obeys

$$('e^{\beta\gamma_5}, \not{p} + e^{i\alpha\gamma_5}m)\psi = 0.$$

This 'free' equation has space-time discrete symmetries:

$$\begin{aligned}\mathcal{P}: \quad & \psi(\vec{x}, t) \rightarrow e^{(\beta-i\alpha)\gamma_5} \gamma_0 \psi(-\vec{x}, t) \\ \mathcal{C}: \quad & \psi(\vec{x}, t) \rightarrow i e^{\beta\gamma_5} \gamma_2 \psi^*(\vec{x}, t) \\ \mathcal{T}: \quad & \psi(\vec{x}, t) \rightarrow -e^{-i\alpha\gamma_5} \gamma_1\gamma_3 \psi^*(-\vec{x}, t)\end{aligned}$$

- Even when theory does not have these symmetries,
- asymptotic states always do,
- but operators have extra γ_5 phases.
- $e^{(-\beta+i\alpha)\gamma_5/2} \psi$ has standard phases.

Dirac Equation

Phase conventions

Consider $N = (\bar{u}^c \gamma_5 d)u$ with usual phases for u and d .

If theory has \mathcal{C} , \mathcal{P} , \mathcal{T} , N has the same phases.

So, in a symmetric theory, $\alpha = \beta = 0$.

Otherwise, by Lorentz invariance, we still have

$$\mathcal{P} = \left[e^{\beta(p^2)\gamma_5} \Pi(p^2) \not{p} - e^{i\alpha(p^2)\gamma_5} \Sigma(p^2) m \right]^{-1}.$$

Like the free Dirac equation, but $e^{(\beta(p^2) - i\alpha(p^2))\gamma_5}$ is not local.

Dirac Equation

State-dependent phases

By Källen-Lehman spectral representation, the propagator is

$$\sum \rho(\mu^2) Z_N(\mu^2) \frac{e^{-\beta(\mu^2)\gamma_5} \not{p} + e^{-i\alpha(\mu^2)\gamma_5} Z_m(\mu^2) \mu}{p^2 - (Z_m(\mu^2) \mu)^2}.$$

When there is no overall symmetry operator, phases state-dependent.

$N_{\text{st}} \equiv e^{(-\beta(m_N^2) + i\alpha(m_N^2))\gamma_5/2} N$ has standard transformations and equation of motion, but only for the neutron; the excited states have non-standard phases.

Dirac Equation

Electric Dipole Moment

$e\sigma \cdot B$ is even under \mathcal{C} , \mathcal{P} and \mathcal{T} ,

$e\sigma \cdot E$ is odd under \mathcal{P} and \mathcal{T} .

$$\Sigma \cdot F \propto \begin{pmatrix} \sigma \cdot B & i\sigma \cdot E \\ i\sigma \cdot E & \sigma \cdot B \end{pmatrix},$$

which is $\sigma \cdot B$ in the rest frame iff $\not{p} \pm m = 0$.

In general, we need to use $e^{i\alpha\gamma_5}\Sigma \cdot F$.

So, important to use N_{st} instead of N in analyses.

At the Green's function level, this is

$$\langle TN_{\text{st}}O\bar{N}_{\text{st}} \rangle = e^{(-\beta(m_N^2) + i\alpha(m_N^2))\gamma_5/2} \langle TNO\bar{N} \rangle e^{(\beta(m_N^2) + i\alpha(m_N^2))\gamma_5/2}.$$

Lattice Calculation

Methods

Two methods for calculating the EDM.



- ① Spin dependent energy in an external electromagnetic field.
 - Need to include background EM fields in gauge generation.
 - Can 'reweight' with disconnected loops.
 - EM flux quantized: so large fields in small volumes.
 - Need derivative at zero.
- ② F_3 Form factor
 - Need to calculate at non-zero momentum transfer.
 - Need derivative at zero-momentum: difficult on small lattices.
 - Need disconnected insertion of currents.
 - Need to account for γ_5 phase.
 - Many operators need 4-point functions, analytic continuation, or source methods.

Lattice Calculation

Topological charge and Weinberg operator

To find the contribution of $\bar{\Theta}$, we need the correlation between the electric current and the topological charge. Similar calculation for the Weinberg operator.

$$\left\langle n \left| \left(\frac{2}{3} \bar{u} \gamma_\mu u - \frac{1}{3} \bar{d} \gamma_\mu d \right) Q \right| n \right\rangle = \frac{1}{2} \left\langle n \left| (\bar{u} \gamma_\mu u + \bar{d} \gamma_\mu d) Q \right| n \right\rangle + \frac{1}{6} \left\langle n \left| (\bar{u} \gamma_\mu u - \bar{d} \gamma_\mu d) Q \right| n \right\rangle$$



Probably no non-zero signal yet.

- Shintani *et al.*, Physical Review **D72** (2005) 014504.
- Berruto *et al.*, Physical Review **D73** (2006) 054509.
- Guo *et al.*, Physical Review Letters **115** (2015) 062001.
- Shindler *et al.*, Physical Review **D92** (2015) 094518.
- Alexandrou *et al.*, Physical Review **D93** (2016) 074503.
- Shintani *et al.*, Physica Review **D93** 094503.

need to be reanalyzed.



Abramczyk estimate that $F_3 \lesssim 0.07\bar{\theta}$ at 1σ .

Lattice Calculation


Quark Electric Dipole Moment

Since the **quark electric dipole moment** directly couples to the electric field, we just need to calculate its matrix elements in the neutron state.


$$\begin{aligned} \langle n | d_u^\gamma \bar{u} \sigma^{\mu\nu} u + d_d^\gamma \bar{d} \sigma^{\mu\nu} d | n \rangle = \\ \frac{d_u^\gamma + d_d^\gamma}{2} \langle n | \bar{u} \sigma^{\mu\nu} u + \bar{d} \sigma^{\mu\nu} d | n \rangle + \frac{d_u^\gamma - d_d^\gamma}{2} \langle n | \bar{u} \sigma^{\mu\nu} u - \bar{d} \sigma^{\mu\nu} d | n \rangle \end{aligned}$$



Parity mixing higher order in α_{EW} : so, result is same as tensor charge.



Results for $u + d$ similar. $g_T^u = 0.792(14)$; $g_T^d = -0.194(14)$.
Disconnected contribution small.




Lattice Calculation

Quark Chromoelectric Moment

nEDM from **quark chromoelectric moment** is a four-point function:

$$\left\langle n \left| \left(\frac{2}{3} \bar{u} \gamma_\mu u - \frac{1}{3} \bar{d} \gamma_\mu d \right) \int d^4x (d_u^G \bar{u} \sigma^{\nu\kappa} u + d_d^G \bar{d} \sigma^{\nu\kappa} d) \tilde{G}_{\nu\kappa} \right| n \right\rangle$$



Lattice Calculation

Reduction to three-point function

The quark chromo-EDM operator is a quark bilinear.

Schwinger source method: Add it to the Dirac operator in the propagator inversion routine:

$$\not{D} + m - \frac{r}{2}D^2 + c_{sw}\Sigma^{\mu\nu}G_{\mu\nu} \longrightarrow \not{D} + m - \frac{r}{2}D^2 + \Sigma^{\mu\nu}(c_{sw}G_{\mu\nu} + i\epsilon\tilde{G}_{\mu\nu})$$


The fermion determinant gives a 'reweighting factor'

$$\begin{aligned} & \frac{\det(\not{D} + m - \frac{r}{2}D^2 + \Sigma^{\mu\nu}(c_{sw}G_{\mu\nu} + i\epsilon\tilde{G}_{\mu\nu}))}{\det(\not{D} + m - \frac{r}{2}D^2 + c_{sw}\Sigma^{\mu\nu}G_{\mu\nu})} \\ &= \exp \text{Tr} \ln \left[1 + i\epsilon \Sigma^{\mu\nu} \tilde{G}_{\mu\nu} (\not{D} + m - \frac{r}{2}D^2 + c_{sw}\Sigma^{\mu\nu}G_{\mu\nu})^{-1} \right] \\ &\approx \exp \left[i\epsilon \text{Tr} \Sigma^{\mu\nu} \tilde{G}_{\mu\nu} (\not{D} + m - \frac{r}{2}D^2 + c_{sw}\Sigma^{\mu\nu}G_{\mu\nu})^{-1} \right]. \end{aligned}$$

Lattice Calculation

Chromo-edm diagrams

$$e^{i\epsilon} \text{ (loop with cross) } \times$$







The chromoEDM operator is dimension 5.

Uncontrolled divergences unless $\epsilon \lesssim 4\pi a \Lambda_{\text{QCD}} \sim 1$.

Need to check linearity.

Two point functions


Neutron Propagator



Preliminary; Connected Diagrams Only

Two point functions

Linearity



Preliminary; Connected Diagrams Only

Use $\epsilon \approx \frac{a}{30\text{fm}} \approx 6.6\text{MeV}$ $a \approx 0.36\text{ma}$ for experiments.



Two point functions

Connected γ_5

$$a(\not{D} + m) + i\epsilon\gamma_5 = e^{\frac{i}{2}\alpha_q\gamma_5} (a\not{D} + am_\epsilon) e^{\frac{i}{2}\alpha_q\gamma_5}$$

where $\alpha_q \equiv \tan^{-1} \frac{\epsilon}{am}$

and $am_\epsilon \equiv \sqrt{(am)^2 + \epsilon^2}$



	a12m310	a09m310
$am^0 \equiv \frac{1}{2\kappa} - 4$	-0.0695	-0.05138
$am_{cr} \equiv \frac{1}{2\kappa_c} - 4$	-0.08058	-0.05943
$am \equiv am^0 - am_{cr}$	0.01108	0.00805
ϵ	0.004	0.003
am_ϵ	0.01178	0.00859
M_π^0	0.1900(4)	0.1404(3)
M_π^{CEDM}	0.1906(4)	0.1407(3)
$M_\pi^{\gamma_5}$	0.1961(4)	0.1450 (3)
$M_\pi^0 \times \sqrt{\frac{m_\epsilon}{m}}$	0.1959(4)	0.1450(3)

Three point functions

F_3 Form factor from CEDM




Figure 4: Signal in F_3 from the insertion of the CEDM operator in the u (left) and d (right) quarks

Preliminary; Connected Diagrams Only

Three point functions

Renormalization

RI-SMOM scheme for non-perturbative renormalization worked out.

Most divergent mixing with $a^{-2}\alpha_s\bar{\psi}\gamma_5\psi$ even with chiral symmetry.
Effect same as of $(\alpha_s/ma^2)G \cdot \tilde{G}$.

Current estimates of nEDM:

- CEDM is $O(1)$.
- $(\alpha_s/ma^2)G \cdot \tilde{G}$ contribution is $O(1)$ - $O(10)$ at $a \sim 0.1\text{fm}$

Not present in connected diagrams with good chiral symmetry!
For Wilson fermions, $O(a)$ chiral breaking gives multiplicative correction.